

Kokusaginina

Punto de fusión, características físicas y UV.

Komala I, Rahmani M, Sukari MA, Ismail HBM, Lian GEC, Rahmat A. 2006. Furoquinoline alkaloids from *Melicope bonwickii* (F Muell) T Hartley. *Nat Resour Res* 20(4): 355-360.

RMN¹H, RMN¹³C y actividad anticolinesterásica.

Cardoso-Lopes EM, Maier JA, Rogério da Silva M, Regasini LO, Simote SY, Lopes NP, Pirani JR, Bolzani VS, Young MCM. 2010. Alkaloids from Stems of *Esenbeckia leiocarpa* Engl. (Rutaceae) as Potential Treatment for Alzheimer Disease. *Molecules* 15: 9205-9213.

EMIE

Orlita A, Sidwa-Gorycka M, Kumirska J, Malinski E, Siedlecka EM, Gajdus J, Lojkowska E, Stepnowski P. 2008. Identification of *Ruta graveolens* L. Metabolites Accumulated in the Presence of Abiotic Elicitors. *Biotechnol Progr* 24(1): 128-133.;

Actividad antipalúdica contra *Plasmodium falciparum*.

Wansi JD, Hussain H, Tcho AT, Kouam SF, Specht S, Sarite SR, Hoerauf A, Krohn K. 2010. Antiplasmodial activities of furoquinoline alkaloids from *Teclea afzelii*. *Phytotherapy Research* 24(5): 775-777.