

## **5-Hidroxi-7,4'-dimetoxiflavona**

### **Características físicas y rayos X**

Kou LQ, Cheng XL, Zhang ZT. 2008. Syntheses and crystal structures of two apigenin alkylation derivatives. *J Chem Crystallogr* 38(1): 21-25.

### **Punto de fusión, UV, IR, RMN<sup>1</sup>H, RMN<sup>13</sup>C y EM ESI-TOF”.**

Sutthanut K, Srpanikulchai B, Yenjai C, Jay M. 2007. Simultaneus identification and quantitation of 11 flavonoid constituents in Kaempferia parviflora by gas chromatography. *J Chromatogr A* 1143(1-2): 227-233.

### **Efecto contra las enzimas α-glucosidasa y α-amilasa**

Krishna MS, Joy B, Sundaresan A. 2015. Effect on oxidative stress, glucose uptake level and lípido droplet content by apigenin 7,4'-dimethyl ether isolated from *Piper longum* L. *J Food Sci Technol* 52(6): 3561-3570.

### **Efecto inhibitorio de citotoxicidad inducida con D-GaIN en hepatocitos de ratón.**

Chaipech S, Morikawa T, Ninomiya K, Yoshikawa M, Pongpiriyadacha Y, Hayakawa T, Muraoka O. 2012. Structures of two new phenolic glycosides, kaempferiaosides A and B, and hepatoprotective constituents from the rhizomes of *Kaempferia parviflora*. *60(1): 62-69.*

### **Actividad antioxidante detectada mediante la técnica teórica CUPRAC, RP and FRAP.**

Ben SM, Skandrami I, Nasr N, Franca MGD, Chekir-Ghedira L, Ghedira K. 2011. Flavonoids and sesquiterpenes from *teucrium ramosissimum* promote antiproliferation of human cancer cells and enhance antioxidant activity: a structure-activity relationship study. *Environ Toxicol Pharmacol* 32(3): 336-348.

### **Otras referencias:**

- Wu T, Jiang C, Wang L, Morris-Natschke SL, Miao H, Gu L, Xu J, Lee KH, Gu Q. 2015. 3,5-Diarypyrazole derivatives obtained by ammonolysis of the total flavonoids from chrysanthemum indicum extract show potencial for the treatment of alzheimer's disease. *J Nat Prod* 78(7): 1593-1599.
- Ganugapati J, Ganesh V, Nasu V. 2015. Virtual screening and Docking Studies of aP2 inhibitors- A key proteína link between asthma and obesity. *J Chem Pharm Res* 7(4): 718-724.
- Li WQ, Zhou MY, Zhou H, Hu XP, Ou SY, Yan RA, Liao XJ, Huang XS, Fu L. 2015. Characterization of phenolic constituents inhibiting the formation of sulfur-containing volatiles produced Turing garlic processing. *J Agric Food Chem* 63(3): 787-794.
- Kim M, Kim N, Han J. 2014. Metabolismo f Kaempferia parviflora polymethoxyflavones by human intestinal bacterium Blautia sp. MRG-PMF1. *J Agric Food Chem* 62(51): 12377-12383.
- Venditti A, Bianco A, Bruno M, Ben JM, Nicoletti M. 2015. Phytochemical study of *Cistus ladanifer* L. *Nat Prod Res* 29(2): 189-192.
- Okabe Y, Shimada T, Horikawa T, Kinoshita K, Koyama K, Ichinose K, Aburada M, Takahashi K. 2014. Supression of adipocyte hypertrophy by polymethoxyflavonoids isolated from Kaempferia parviflora. *Phytomedicine* 21(6): 800-806.
- Coulierie P, Nour M, Maciuk A, Eydoux C, Gullemot JC, Lebouvier N, Hnawia E, Leblanc K, Lewin G, Canard B, et al. 2013. Structure-activity relationship study of biflavonoids on the dengue virus polymerase DENV-NS5 RdRp. *Planta Med* 79(14): 1313-1318.
- Sasaki H, Kitoh Y, Miki K, Kinoshita K, Koyama K, Kaneda M, Takahashi K. 2012. Structure-relationships of biflavonoids for b-secretase (BACE-1) inhibitory activity. *Heterocycles* 85(11): 2749-2756.

- Liu Y, Xu XH, Liu Z, Du XL, Chen KH, Xin X, Jin ZD, Shen JZ, Hu Y, Li GR, et al. 2012. Effects of the natural flavone trimethylapigenin on cardiac potassium currents. *Biochem Pharmacol* 84(4): 498-506.
- Sae-Wong C, Matsuda H, Tewtrakul S, Tansakul P, Nakamura S, Nomura Y, Yoshikawa M. 2011. Suppressive effects of methoxyflavonoids isolated from *Kaempferia parviflora* on inducible nitric oxide synthase (iNOS) expresión in raw 264.7 cells. *J Ethnopharmacol* 136(3): 488-495.
- Vera N, Solorzano E, Ordonez R, Maldonado L, Bedascarrasbure E, Isla MI. 2011. Chemical composition of Argentinian propolis collected in extreme regions and its relation with antimicrobial and antioxidant activities. *Nat Prod Commun* 6(6): 823-827.
- Savi LA, Caon T, De Oliveira AP, Sobottaka AM, Werner W, Reginatto FH, Schenkel EP, Barardi CRM, Simoes CMO. 2010. Evaluation of antirotavirus activity of flavonoids. *Fitoterapia* 81(8): 1142-1146.
- Sawasdee P, Sabaphon C, Sitthiwonwanit D, Kokpol U. 2009. Anticholinesterase activity of 7-methoxyflavones isolated from *Kaempferia parviflora*. *Phytoterapy Res* 23(12): 1792-1794.
- Wollenweber E, Fischer R, Doerr M, Irvine K, Pereira C, Stevens JF. 2008. Chemodiversity of exudate flavonoids in *Cassinia* and *Ozothamnus* (Asteraceae, Gnaphalieae). *Z Naturforsch C: Biosci* 63(9/10): 731-739.
- Tewtrakul S, Subhadhirasakul S. 2008. Effects of compounds from *Kaempferia parviflora* on nitric oxide, prostaglandina E2 and tumor necrosis factor-alpha productions in RAW264.7 macrophage cells. *J Ethnopharmacol* 120(1): 81-84.
- Zhao J, Pawar RS, Ali Z, Khan IA. 2007. Phytochemical investigation of *Turnera diffusa*. *J Nat Prod* 70(2): 289-292.
- Yilmaz BS, Citoglu GS, 2006. High performance liquid chromatography liquid chromatographic analysis of some flavonoids of *Ballota* species. *Chem Nat Prod* 42(3): 353-355.
- Nikolova MT, Taskova RM, Peev DR. 2005. Exudate flavonoid aglycones of *Veronica*: ecological and systematic implications. *Biochem Syst Ecol* 33(12): 1258-1268.
- Oluwatuy M, Kaatz GW, Gibbons S. 2004. Antibacterial and resistance modifying activity of *Rosmarinus officinalis*. *Phytochemistry* 65(24): 3249-3254.
- Topcu G, Altiner EN, Gozcu S, Helfon B, Aydogmus Z, Pezzuto JM, Zhou BN, Kingston DGI. 2003. Studies on Di- and triterpenoids from *Salvia staminea* with cytotoxic activity. *Planta Med* 69(5): 464-467.
- Karzhaubekova ZZ, Burasheva GS. 2002. Phytochemical study of *Kalidium caspicum*. *Chem Nat Compd* 38(1): 100-101.
- Grayer RJ, Veitch NC, Kite GC, Price AM, Kokubun T. 2001. Distribution of 8-oxygenated leaf-surface flavones in the genus *Ocimum*. *Phytochemistry* 56(6): 559-567.
- Ortega A, Mondragon P, Maldonado E. 1999. Guaianolides of *Stevia laxiflora*. *Rev Soc Quim Mex* 43(3,4): 100-102.
- Quijano L, Trejo RI, Collera O, Rios T. 1994. Sesquiterpene lactones from *Montanoa leucantha* subsp. *leucantha*. *Phytochemistry* 36(6): 1443-1448.
- Jacobs H, Bunbury M, Hylands PJ, McLean S. 1986. Flavonoids and a heliangolide, nivesuin C, from *Melampodium camphoratum*. *J Nat Prod* 49(6): 1163-1164.
- Edwards JM, Raffauf RF, Le Quesne PW. 1979. Antitumor plants. Part VII. Antineoplastic activity and cytotoxicity of flacones, isoflavones, and flavanones. *J Nat Prod* 42(1): 85-91.
- Nogradi M, Farkas L, Wagner H, Hoerhammer L. 1967. Transacylation reaction in the flavonoid series. II. New synthesis of cosmosiine and tilianin. *Ber* 100(9): 2783-2790.
- Hasegawa M, Shirato T. 1955. Flavonoids of various *Prunus* species. IV. The flavonoids in the wood of *Prunus donarium* var. *spontanea*. *J Am Chem Soc* 77, 3557-3558.
- Bauer KH, Dietrich H. 1933. The pigment of birch buds. *Berichte der Deutschen Chemischen Gesellschaft B: Abhandlungen* 66B: 1053-1054.
- Walz E. 1931. Isoflavone and saponina glucosides in *Soja hispida*. *Justus Liebigs Annalen der Chemie* 489: 118-155.

