

Geraniol

RMN¹H

Howell AR, Pattenden G. 1990. Hydrocobaltation reactions of 1,3-dienes. Regioselective hydroxylation of myrcene to geraniol and to (±)-linalool via allylcobaloxime intermediates. J Chem Soc Perkin Trans 1: (10): 2715-2720.

EMIE, P.E.

Won MM, Cha EJ, Yoon OK, Kim NS, Kim K, Lee DS. 2009. Use of headspace mulberry paper bag micro solid phase extraction for characterization of volatile aromas of essential oils from Bulgarian rose and Provence lavender. Anal Chim Acta 631(1): 54-61.

Actividad moderada LC₅₀ > 10 mg/cm² contra *Blattella germánica*

Jang YS, Yang YC, Choi DS, Ahn YJ. 2005. Vapor phase toxicity of marjoram oil compounds and their related monoterpenoids to *Blattella germanica* (Orthoptera: Blattellidae). J Agric Food Chem 53(20): 7892-7898.

CMI 200 y CML 250 ppm contra *Saprolegnia parasítica*

Tampieri MP, Galuppi R, Carelle MS, Macchioni F, Cioni PL, Morelli I. 2003. Effect of selected essential oils and pure compounds on *Saprolegnia parasitica*. Pharm Biol 41(8): 584–591.