

7,4'-Dimetoxi naringenina

UV, IR, EMIE, RMN¹H y RMN¹³C.

Wesolowska O, Wisniewski J, Duarte N, Ferreira MJU, Michalak K. 2007. Inhibition of MRP1 transport activity by phenolic and terpenic compounds isolated from Euphorbia species. *Anticancer Res* 27(6B): 4127-4134.

Punto de fusión

Rossi MH, Yoshida M, Maia JGS. 1997. Neolignans, styrylpyrones and flavonoids from Aniba species. *Phytochemistry* 45(6): 1263-1269.

Presentó actividad antifúngica frente a *Cladosporium cladosporioides* y *Cladosporium sphaerospermum*

Lago JHG, Young MCM, Reigada JB, Soares MG, Roesler BP, Kato MJ. 2007. Antifungal derivatives from Piper mollicomum and P. Ihotzkyanum (Piperaceae). *Quim Nova* 30(5): 1222-1224.

Otras referencias

- McDonald BR, Nibbs AE, Scheidt KA. 2015. A biomimetic strategy to Access the silybins: total synthesis of (-)-isosilybin A. *Organic Lett* 17(1): 98-101.
- Fareza MS, Syah YM, Mujahidin D, Juliawaty LD, Kurniasih I. 2014. Antibacterial flavanone and dihydrochalcones from Macaranga trichocarpa. *Z Naturforsch C: Biosci* 69(9/10): 375-380.
- Lu K, Chu J, Wang H, Fu X, Quan D, Ding H, Yao Q, Yu P. 2013. Regioselective iodination of flavonoids by N-iodosuccinimide under neutral conditions. *Tetrahedron Lett* 54(47): 6345-6348.
- Zhang NL, Zhu YH, Huang RM, Fu MQ, Su ZW, Cai JZ, Hu YJ, Qiu SX. 2012. Two new stilbenoids from Cajanus cajan. *Z Naturforsch B: Chemical Sci* 67(12): 1314-1318.
- Deng Y, Chin YW, Chai HB, Carcache de Blanco E, Kardono LBS, Riswan S, Soejarto DD, Farnsworth NR, Kinghorn AD. 2011. Phytochemical and bioactivity studies on constituents of the leaves of Vitex quinata. *Phytochemistry Lett* 4(3): 213-217.
- Cao CM, Xu LJ, Peng Y, Shi QW, Xiao PG. 2010. Two new flavan-flavanones from Sarcandra hainanensis. *Chem Pharm Bull* 58(10): 1395-1398.
- Wollenweber E, Wehde R, Christ M, Doerr M. 2008. Surface flavonoids in Catalpa ovata, Greyia sutherlandii and Paulownia tomentosa. *Nat Prod Commun* 3(8): 1285-1287.
- Pisutthanan N, Liawruangrath B, Liawruangrath S, Bremner JB. 2006. A new flavonoid from Chromolaena odorata. *Nat Prod Res* 20(13): 1192-1198.
- Cai XH, Luo XD, Zhou J, Hao XJ. 2005. Dolabellane diterpenoids from the higher plant Aglaia odorata. *Helv Chim Acta* 88(11): 2938-2943.
- Wollenweber E, Doerr M, Bohm BA, Roitman JN. 2004. Exudate flavonoids of eight species of Ceanothus (Rhamnaceae). *Z Naturforsch C: Biosci* 59(7/8): 459-462.
- Miyazawa M, Kinoshita H, Okuno Y. 2003. Antimutagenic activity of sakuranetina from Prunus jamasakura. *J Food Sci* 68(1): 52-56.
- Kamperdick C, Hong Van N, Van Sung T. 2002. Constituents from Miliusa balansae (Annonaceae). *Phytochemistry* 61(8): 991-994.
- Schmidt TJ, Willuhn G. 2000. Sesquiterpene lactone and flavonoid variability of the Arnica angustifolia aggregate (Asteraceae). *Biochem Syst Ecol* 28(2): 133-142.
- Wollenweber E, Doerr M, Muniappan R. 1995. Exudate flavonoids in a tropical weed, Chromolaena odorata (L.) R. M. King et H. Robinson. *Biochem Syst Ecol* 23(7/8): 873-874.
- Sanz JF, Garcia-Lliso V, Marco JA, Valles-Xirau J. 1991. A cadinane derivative from Artemisia crithmifolia. *Phytochemistry* 30(12): 4167-4168.

- Echeverri F, Suarez GE. 1989. Flavonoids from *Passiflora foetida* and deterrent activity. *Rev Latinoam Quim* 20(1): 6-7.
- Bohlmann F, Scheidges C, Zdero C, King RM, Robinson H. 1984. ent-Labdanes from *Baccharis sternbergiana*. *Phytochemistry* 23(5): 1109-1111.
- Bohlmann F, Zdero C, King RM, Robinson H. 1980. Naturally occurring terpene derivative. Part 289. Seven guaianolides from the tribe Vernoniae. *Phytochemistry* 19(12): 2669-2673.
- Lam J, Wrang P. 1975. Chemical constituents of the genus *Dahlia*. 6. Flavonoids and polyacetylenes in *Dahlia tenuicaulis*. 1975. *Phytochemistry* 14(7): 1621-1623.
- Bohm BA. 1968. Phenolic compounds in ferns. II. Indirect evidence for the existence of 2',6'-dihydroxy-4,4'-dimethoxychalcone in *Pityrogramma calomelanos*. *Phytochemistry* 7(9): 1687-1689.